

# ANNALS OF PHYSICAL MEDICINE AND PHYSIOTHERAPY

## Effect of Low-Level Laser Therapy and Manual Exercises on Shoulder Impingement Syndrome: A Systematic Review

El-Gendy MH<sup>1</sup>,  
Abdulla YM<sup>2</sup>,  
Abdalla GAE<sup>3</sup> and  
Elmelgie MM<sup>4\*</sup>

<sup>1</sup>Department of Physical Therapy for Basic Sciences, Faculty of Physical Therapy, Cairo University

<sup>2</sup>Department of Physical Therapy for Basic Sciences, Faculty of Physical Therapy, Cairo University

<sup>3</sup>Assistant Professor, Department of Physical Therapy for Basic Sciences, Faculty of Physical Therapy, Cairo University

<sup>4</sup>Lecturer, Department of Physical Therapy for Basic Sciences, Faculty of Physical Therapy, October 6 University

### Article Information

<b>Article Type:</b>	Review Article	<b>*Corresponding Author:</b>	<b>Citation:</b>
<b>Journal Type:</b>	Open Access	Mohamed M. ElMelgie,	Elmelgie MH (2021). Effect of Low-Level Laser Therapy and Manual Exercises on Shoulder Impingement Syndrome: A Systematic Review. Ann Phys Med Physiother. 1(1); 1-7
<b>Volume:</b>	Issue: 1	Lecturer, Department of Physical Therapy for Basic Sciences, Faculty of Physical Therapy, October 6 University,	
<b>Manuscript ID:</b>	APMP-v1-101	E-mail: Mohamed.magdy.pt@o6u.edu.eg	
<b>Publisher:</b>	Science World Publishing		
<b>Received Date:</b>	10 May 2021		
<b>Accepted Date:</b>	31 May 2021		
<b>Published Date:</b>	11 June 2021		

**Copyright:** © 2021, Elmelgie MH, *et al.*, This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 international License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

### ABSTRACT

**BACKGROUND:** The aim of this work was to systematically review the influence of low level laser therapy and manual exercises used on the patients with shoulder impingement syndrome.

**METHODS:** Systematic review of all published studies with all research designs except expert opinions. A search was made in Pubmed, Medline, Cochrane library, PEDro and Google scholar; from the earliest date to September 2019.

**INTERVENTION:** Low level laser therapy and manual exercises programs performed by the physical therapist. **RESULTS:** Only 4 studies met the inclusion criteria. Meta-analysis could be done and findings are presented qualitatively due to heterogeneity of the studies. There is no conflicting evidence on whether low level laser therapy and manual exercises can increase the range of movement and decrease pain, for those studies showing improvements in the range of movements.

**CONCLUSION:** The current level of evidence support the effectiveness of low level laser therapy and manual exercises in patients with shoulder impingement syndrome remains.

**KEYWORDS:** Low level laser therapy, Manual exercises, Shoulder impingement syndrome, Systematic Review.

### INTRODUCTION

Shoulder Impingement Syndrome (SIS) occurs due to a mechanical disturbance within the subacromial space and is characterized by pain and functional restrictions mostly during overhead activities in daily life or sporting activities [1].

Potential factors causing or contributing to SIS such as strength, coordination and integrity of the rotator cuff [2] and the shoulder girdle muscles [3] mechanical or anatomical changes, hypomobility or instability of the glenohumeral joint or the scapula, and the influence of posture [4] are discussed in the literature and suggest a multi-factorial etiology of SIS. Besides the biomedical aspects of SIS, psychological factors such as kinesiphobia or catastrophizing may negatively influence recovery and thus leading to chronic pain and disability [5].

The specific diagnosis of SIS is often based on a thorough history and clinical examination; technical examination methods such as MRI or ultrasonography are often not used in first instance [6],

also because their diagnostic accuracy is still limited [7].

Physiotherapy is often the first choice of treatment for SIS. Between 10 to 30% of all shoulder patients seen in primary care are referred to physiotherapy after initial presentation [6]. However, the effectiveness of physiotherapy in patients with SIS is still under debate. Conclusions from systematic reviews suggest that physiotherapy-led interventions, combining different methods or techniques, are not more effective than exercises alone except adding manual mobilization to exercises, which seems to be of additional benefit. Most technical treatments such as ultrasound or laser therapy cannot be recommended. However, evidence is limited by poor methodological quality, short follow ups and small sample sizes [8].

Treatment for it divided into conservative treatment and surgical treatment. Conservative treatment involves rest, medication, exercise, and electrotherapy. In early-stage impingement syndrome, conservative treatment is effective. In general, it is desirable to perform exercise treatment after the pain has subsided and the soft tissues have been relaxed by a warming and cooling treatment and electrotherapy [9].

A comprehensive manual physical therapy approach, including both thrust and non-thrust techniques and reinforcing mobility exercises, is a common intervention for patients presenting with a primary report of shoulder pain. This comprehensive approach has shown improvement in shoulder symptoms with mobilisation and manipulation techniques targeted to the thoracic spine, cervicothoracic spine, rib cage and acromioclavicular joints, in addition to the glenohumeral joint [10].

Laser is a noninvasive, nonionising, monochromatic electromagnetic high concentrated light beam. Recently, Low Level Laser Therapy (LLLT) is widely used in various rheumatologic and musculoskeletal disorders which have analgesic, anti-inflammatory and biostimulating effects. The LLLT induces cell proliferation, collagen synthesis, protein synthesis, tissue reparation, wound healing and pain relief through direct irradiation without thermal response [11].

However, conflicting results were reported about the effectiveness of LLLT on musculoskeletal disorders. Some of the randomized controlled studies suggest that LLLT may be effective in pain relief in different musculoskeletal disorders [12]. On the other hand some of them have failed to show any superiority over placebo [13].

## **SUBJECT AND METHODS**

### **SEARCH STRATEGY FOR IDENTIFICATION OF STUDIES:**

Electronic database search was performed from the earliest date.

The study was applied between January to December 2020 (Table 1) to identify relevant articles in:

- PubMed (Medline) at <http://www.ncbi.nlm.nih.gov/pubmed>
- The Cochrane Library at <http://www.thecochranelibrary.com>
- Physiotherapy Evidence Database (PEDro) at <http://www.pedro.org.au/>
- Google scholar at <http://scholar.google.com>

The following key words were used in the search:

- "Subacromial impingement syndrome".
- "Shoulder impingement syndrome".
- "Low level laser therapy"
- "Physical therapy exercises".

Reference lists in the relevant studies and review articles were examined.

### **STUDY SELECTION CRITERIA:**

The titles and abstracts collected by the above mentioned search strategy, were initially screened against the inclusion and exclusion criteria for identification of the relevant trials. When the title and abstract did not indicate clearly if an article should be included, the complete article would be read to determine its suitability.

### **INCLUSION CRITERIA:**

#### **TYPES OF STUDIES:**

Published full text articles in peer-reviewed journals with all research designs except expert opinions.

#### **TYPES OF PARTICIPANTS:**

The review included children (from 20 to 55 years of age).

#### **TYPES OF INTERVENTIONS:**

This review included studies which demonstrate the effects of passive manual stretching programs performed by the physical therapist with reported findings for analysis of its effectiveness.

#### **TYPES OF OUTCOME MEASURES:**

Only outcome measures related to passive joint range of motion were considered in this review.

### **EXCLUSION CRITERIA**

#### **UNPUBLISHED STUDIES:**

- Studies that compared passive stretching programs with the effects of medications, surgery, or serial casting were excluded as the area of interest was mainly on passive stretching without assistance from surgery and antispasticity medications.
- Studies that measured the effect of stretching on spastici-

ty or gait parameters.

- Studies that combined stretching with other types of modalities; such as heating, therapeutic ultrasound, splinting and electrical stimulation.

#### **QUALITY ASSESSMENT OF METHODOLOGY:**

All the included studies were scored on their methodological rigour with the Physiotherapy Evidence Database (PEDro) scale (PEDro, 2010).

#### **EXPLANATION OF SOME RESEARCH DESIGNS:**

##### **N-OF-1 RANDOMIZED CONTROLLED TRIALS:**

In single subject research, treatment versus control conditions are manipulated within a single person; the order of these exposures is randomly allocated. There are several variations of the N-of-1 RCT, sometimes called a randomized cross-over trial; these include the blind cross-over trial or double blind cross-over trial.

The difference between this and a group crossover is that there are repeated measures in multiple phases. A person frequently undergoes pairs of periods in which one period applies an experimental treatment (B) and the other applies a placebo (C) or baseline (A)-in other words, an ABABA type of design or ABCBCBA or variation. The order of these periods within each pair is randomly selected so that the conduct of the trial may be, for example, ABBAAB.

Treatment outcomes are monitored to document the effect of the condition currently being applied. These phases are repeatedly measured until the person being treated and the investigator are convinced that the treatment period is clearly different, or clearly not different. In a blind trial, the person making the outcome assessments is blind to the treatment condition; in a double blind trial, both the subject and the assessor are unaware of the treatment condition.

Although this method can also provide a group comparison when more than one subject has been studied, the focus of the published report is the individual comparisons. Alternatively, when multiple N-of-1 randomized controlled trials conducted under the same protocol have been summed and a group comparison is provided, this is called a multiple cross-over trial.

Another variation of the N-of-1 RCT is the alternating treatments design in which the subject is exposed to the treatment condition and control condition(s) in close temporal proximity. For example, a subject is assessed during a 20minute exposure to a control condition followed by a 20 minute exposure to the treatment condition; these exposures are determined by random allocation. Yet another variation is the multiple baseline across subjects design; several subjects are assessed for differing periods of exposure to the nontreatment condition (called baseline) and then assessed

during treatment exposure. The order in which subjects change from the control condition to the treatment condition is established through random allocation.

#### **RESULTS**

Only four studies met the inclusion criteria. Randomized controlled trials were made on the topic. The main reasons for exclusion of the other studies were:

- The other study doesn't meet the inclusion criteria.
- Other types of sample stratified as convenient samples.
- Published articles in non-English language.

#### **METHODOLOGICAL QUALITY RESULTS:**

The scoring of each study with the Physiotherapy Evidence Database (PEDro) scale is listed in (Table 2). The scores of the all studies included in the study ranges from six to seven, the more the number of scores of the aspects evaluating the quality of the study, the more quality of the study.

There are four studies (Table 3) investigating the influence of low level laser therapy and exercises use on the patients with shoulder impingement syndrome.

The study by Dogan et al., (2010) of level II evidence showed that there was in group I a statistically significant improvements in pain severity, range of motion except internal and external rotation and SPADI scores were observed compared to baseline scores after the therapy ( $p<0.05$ ). In group II, all parameters except range of motion of external rotation were improved ( $p<0.05$ ). However, no significant differences were recorded between the groups ( $p<0.05$ ).

The study by Kelle and Kozanoglu, (2014) of level I evidence showed that there was a significant differences were observed between groups I and II and between groups II and III regarding pain during activity and at rest scores at all of the visits ( $p<0.05$ ). Nevertheless, significant improvement was observed between groups I and III regarding pain during activity only at post-treatment ( $p=0.013$ ). The UCLA scores were significantly changed in all three study groups at all of the visits ( $p<0.05$ ).

The study by Pinar et al., (2015) of level II evidence showed that there was a statistically significant improvement in all outcome measurements in both groups. In comparison of both groups, median active and passive flexion, active and passive abduction, passive internal rotation, and passive external rotation scores were significantly higher in the laser group after the treatment ( $p=0.015$ ,  $P=0.004$ ,  $p=0.048$ ,  $p=0.014$ ,  $p=0.031$ , and  $p=0.044$ , respectively), but median differences in both groups were similar ( $P>0.05$ ). The median differences for global visual analogue scale pain and DASH-T scores, in both groups were also similar ( $P>0.05$ ).

**Table 1:** Search results

	Search strategy	PubMed results	Cochrane results	PEDro results	Google results
1	"Subacromial impingement syndrome" AND "Shoulder impingement syndrome"	152	23	9	324
2	"Shoulder impingement syndrome" AND "Low level laser therapy"	121	19	5	198
3	"Subacromial impingement syndrome" AND "Physical therapy exercises"	205	24	4	324
4	"Low level laser therapy" AND "Physical therapy exercises"	102	23	1	201
5	All Keywords combined	1	1	0	99

**Table 2:** Methodology assessment of studies according to the Physiotherapy Evidence Database (PEDro) scale.

Criteria	Dogan et al., (2010).	Thornton et al., (2013).	Kelle and Kozanoglu (2014).	Pinar et al., (2015).	Awotidebe et al., (2015).
1-Specified eligibility criteria	Yes	Yes	Yes	Yes	Yes
2-Random allocation of participants	Yes	Yes	Yes	Yes	Yes
3-Concealed allocation	Yes	No	No	No	No
4-Similar prognosis at baseline	Yes	Yes	Yes	Yes	Yes
5-Blinded participant	No	No	No	No	Yes
6-Blinded therapists	No	Yes	No	No	No
7-Blinded assessors	No	Yes	No	Yes	No
8-More than 85% follow-up for at least one key outcome	Yes	Yes	Yes	No	Yes
9-Intention to treat' analysis	No	No	Yes	No	Yes
10-Between group statistical analysis for at least one key outcome	No	Yes	Yes	Yes	Yes
11-Point estimates of variability for at least one key out come	Yes	Yes	No	Yes	Yes
PEDro score	10-Jun	10-Aug	10-Jun	10-Jun	10-Aug

**Table 3:** Summarizes the characteristics of the research participants in these four studies.

	Dogan et al., (2010).	Kelle and Kozanoglu (2014).	Pinar et al., (2015).	Awotidebe et al., (2015).
Research design	-Multiple single-subject design (a randomized placebo controlled double-blind prospective study).	-A controlled clinical trial.	-A randomized, placebo controlled, prospective study	-A systematic review protocol
Level of evidence.	II	I	II	II
Participants' characteristics.	-Subacromial impingement syndrome	-Subacromial impingement syndrome.	-Subacromial impingement syndrome.	-Subacromial impingement syndrome.
No. participants.	52	135	60	70
Age.	45-60	30-45	18-75	18-35
Intervention.	-Low level laser therapy and placebo laser therapy	-Local corticosteroid injection, sham laser treatment and low-level laser treatment.	-Low level laser therapy and placebo laser therapy.	-Low-level laser therapy with exercises and exercises alone.
Control intervention.	-Improvements in pain severity and range of motion.	-Improvements in pain severity and range of motion.	-Improvements in pain severity and range of motion.	-Improvements in pain severity and range of motion.
Outcome of interest.	-Activities of daily living.	-Activities of daily living.	-Activities of daily living.	-Activities of daily living.

<b>Measures.</b>	-Pain severity. - Range of motion. - Functional status.	-Visual analog scale (VAS) during activity and at rest. - Shoulder functional status and quality of life measured by the University of California at Los Angeles rating score (UCLA). - Nottingham Health Profile (NHP) scale respectively.	-Global visual analogue scale pain. - DASH-T scores.	-Visual analogue scale (VAS) and other categorical rating scales )the higher the rating, the higher the pain).
<b>Component of health.</b>	-Activity and participation.	-Activity and participation.	-Activity and participation.	-Activity and participation.
<b>Results.</b>	-Statistically significant improvements in pain severity, range of motion except internal and external rotation (group I) - All parameters except range of motion of external rotation were improved (group II).	-Significant differences were observed between groups I and II and between groups II and III regarding pain during activity and at rest scores at all of the visits. - Nevertheless, significant improvement was observed between groups I and III regarding pain during activity only at post-treatment. - The UCLA scores were significantly changed in all three study groups at all of the visits.	-Improvements in all outcome measurements in both groups. - In comparison of both groups, median active and passive flexion, active and passive abduction, passive internal rotation, and passive external rotation scores were significantly higher in the laser group after the treatment.	-Significant improvement in all measured variables in both groups (low level laser therapy and exercises group and exercises only group) in favor of low level laser therapy and exercises group.

The study by Awotidebe et al., (2015) of level II evidence showed that there was a statistically significant improvement in all outcome measurements in both groups. Both Chi2 test and I2 statistic will be interpreted based on the guidelines recommended by the Cochrane Handbook for Systematic Review of Interventions ( $P>0.05$ ).

## DISCUSSION

The low level laser therapy has become popular in the treatment of musculoskeletal disorders in recent years. The mechanism of analgesic effect of LLLT is not well known. LLLT can modulate inflammatory pain by reducing levels of biochemical markers, neutrophil cell influx, oxidative stress, and the formation of edema in a dose-dependent manner. The other mechanisms suggested concerning the pain relieving effect of LLLT are altering excitation and nerve conduction in peripheral nerves and stimulation the release of endogenous endorphins [14].

LLLT may accelerate collateral circulation and enhance microcirculation so as to normalize the functional features of the injured

areas. LLLT may also reduce histological abnormalities, collagen concentration, and oxidative stress [15]. LLLT is suggested to be able to accelerate the healing process of tendinous tissue after an injury, increasing fibroblast cell proliferation and collagen synthesis. The clinical effectiveness of LLLT is debatable because of the lack of consensus about the dosage to be used, delivery system, and the wavelength to be delivered [16].

Also, [17] reported that LLLT combined with exercise was more effective than exercise therapy alone in pain relief and increased shoulder ROM. In a systematic review, [18] indicated that exercise therapy was effective in recovery from rotator cuff disease, but laser therapy was not effective for rotator cuff tendinitis.

Because of the additive effect of co-interventions, an isolated effect of LLLT might have not been elucidated. In addition, [19] pointed out that laser therapy was effective only when used in isolation and not when combined with exercise. Laser therapy as a single intervention was recommended in patients who were unable to perform therapeutic exercises.



The results of the clinical trials about treatment of musculoskeletal pain syndromes with LLLT seem controversial. [20] reported in a meta-analysis that the trials supporting the positive effect of LLLT had higher methodological quality. On the other hand in another metaanalysis, [21] suggested that efficacy of LLLT was found to be lower in double blind trials when compared with the uncontrolled ones. This inconsistency is also valid for the effect of LLLT in the therapy of MPS.

In a study, [22] suggested that LLLT had an effect on the trigger points and that the treatment significantly increased the pain threshold while [23] reported no beneficial effect over placebo in a controlled cross-over study during 5 weeks follow up. Why the effect of LLLT is controversial on trigger points may be due to many reasons. Methodological differences in patient selection, trigger points treated (active or inactive) outcome measures selected, and the application parameters of LLLT (wavelength, intensity, duration) may affect the final improvement in pain or functional limitation.

The LLLT has a positive effect on acute and chronic musculoskeletal pain. Due to the heterogeneity of populations, interventions and comparison groups, this diversity means that every single study has not been positive. Pain is a very complex condition which presents in different forms with an interplay of mechanical, biochemical, psychological and socioeconomic factors [24].

It is extremely challenging to compare LLLT to other treatments, and LLLT regimens are complicated by different lengths of treatment, all without standardization of wavelengths and dosages. There have been no long-term (greater than 2 year follow up) human clinical studies that have evaluated LLLT. The overall positive short term clinical studies in addition to strong laboratory studies should give the clinical confidence that LLLT may be beneficial for many individuals suffering from musculoskeletal pain, regardless of the cause [25].

Manual therapy has been shown to be effective at augmenting the effect of exercise in relieving symptoms of the impingement syndrome. Manual therapy includes a variety of techniques, including joint mobilization, and soft tissue mobilization (effleurage, friction, and kneading techniques) [26].

Different studies have determined that manual and exercise therapy appear to have a role in shoulder impingement [27]. It is observed that a 4-week program along with motor management and strengthening exercises reduced shoulder pain and improved feature of individuals with shoulder impingement syndrome. The efficacy of ultrasound, laser, and exercises of shoulder impingement syndrome has been emphasized [28].

Additionally, another study suggested that manual therapy, combined with supervised exercise therapy, alleviated SAIS by means

of decreasing pain and improving shoulder muscle strength and functionality [29]. It has been proposed that manual therapy and stretching exercises aimed at the glenohumeral joint (posteroinferior soft tissue structures) and thoracic spine, seemed to be more effective in the majority of SAIS patients [30].

## CONCLUSION:

Low level laser therapy and manual exercises use on the patients with shoulder impingement syndrome and significant effect on the activities of their daily living with maintenance of this effect for a long period of time.

## REFERENCES

1. Lewis JS, Green AS, Dekel SP. "The aetiology of subacromial impingement syndrome". *Physiotherapy*. 2001; 87: 458-69.
2. Irlenbusch U, Ganssen H. "Muscle biopsy investigations on neuromuscular insufficiency of the rotator cuff: a contribution to the functional impingement syndrome of the shoulder". *J Shoulder Elbow Surg*. 2003; 12: 422-26.
3. Cools A, Dewitte V, Lanszweert F, Notebaert D, Roets A, Soetens B, et al. Rehabilitation of scapular muscle balance: which exercises to prescribe?. *Am J Sports Med*. 2007; 35: 1744-50.
4. Bullock MP, Foster NE, Wright CC. "Shoulder impingement: the effect of sitting posture on shoulder pain and range of motion". *Man Ther*. 2005; 10: 28-37.
5. Feleus A, Van Dalen T, Bierma-Zeinstra S, Bernsen R, Verhaar J, Koes B, et al. "Kinesiophobia in patients with non traumatic arm, neck and shoulder complaints: a prospective cohort study in general practice". *BMC Musculoskeletal Disorders*. 2007; 8: 117-23.
6. Linsell L, Dawson J, Zondervan K, Rose P, Randall T, Fitzpatrick R Carr A. "Prevalence and incidence of adults consulting for shoulder conditions in UK primary care: patterns of diagnosis and referral". *Rheumatology*. 2006; 45: 215-21.
7. Awerbuch MS. "The clinical utility of ultrasonography for rotator cuff disease, shoulder impingement syndrome and subacromial bursitis". *Med J Aust*. 2008; 188: 50-3.
8. Kuhn JE. "Exercise in the treatment of rotator cuff impingement: A systematic review and a synthesized evidence-based rehabilitation protocol". *J Shoulder Elbow Surg*. 2009; 18: 138-60.
9. Park JY, Kim MH. "Changes of ranges of motion according to ages and manifestation frequency of impingement sign in shoulder impingement syndrome". *J Korean Orthop Assoc*. 2007; 32: 384-90.
10. Rhon DI, Boyles RE, Cleland JA, Brown DL. "A manual physical therapy approach versus subacromial corticosteroid injection for treatment of shoulder impingement syndrome: a protocol for a randomised clinical trial". *BMJ Open accessible medical research*. 2011; 1- 12.
11. Dogan SK, Saime AY, Evcik DI. "The effectiveness of low laser therapy in subacromial impingement syndrome: a randomized placebo

- controlled double-blind prospective study". *CLINICAL SCIENCE*. 2010; 65: 1019-22.
12. Basford JR, Sheffield CG, Harmsen WS. "Laser therapy: a randomized, controlled trial of the effects of low-intensity Nd: YAG laser irradiation on musculoskeletal back pain". *Arch Phys Med Rehabil*. 1999; 80: 647-52.
  13. Kulekcioglu S, Sivrioglu K, Ozcan O, Parlak M. "Effectiveness of low-level laser therapy in temporomandibular disorder". *Scand J Rheumatol*. 2003; 32: 114-8.
  14. Ihsan FR. "Low-level laser therapy accelerates collateral circulation and enhances microcirculation". *Photomed Laser Surg*. 2005; 23: 289-94.
  15. Carrinho PM, Renno AC, Koeke PM, Salate AC, Parizotto NA, Vidal BC. "Comparative study using 685-nm and 830-nm lasers in the tissue repair of tenotomized tendons in the mouse". *Photomed Laser Surg*. 2006; 24: 754-8.
  16. Montes-Molina R, Martínez-Rodríguez M, Rodríguez A, Martínez-Ruiz F, Prieto-Baquero A. "Interferential light therapy in the treatment of shoulder tendinopathies: A randomized controlled pilot study". *Clin. Rehabil*. 2012; 26: 1114-22.
  17. Abrisham S, Kermani-Alghoraishi M, Ghahramani R, Jabbari L, Jomeh H, Zare M. "Additive effects of low-level laser therapy with exercise on subacromial syndrome: A randomized, double-blind, controlled trial". *Clin Rheumatol*. 2011; 30: 1341-6.
  18. Green S, Buchbinder R, Hetrick S. "Physiotherapy interventions for shoulder pain". *Cochrane Database Syst Rev*. 2: CD004258. 2003.
  19. Michener LA, McClure PW, Karduna AR. "Anatomical and biomechanical mechanisms of subacromial impingement syndrome". *Clin Biomech (Bristol, Avon)*. 2003; 18: 369-79.
  20. Beckerman H, Bie R, Bouter L, Cuyper H, Oostendorp RA. "The efficacy of laser therapy for musculoskeletal and skin disorders: A criteria based meta-analysis of randomised clinical trials". *Phys Ther*. 2002; 72: 483-91.
  21. Gam A, Thorsen H, Lonnberg F. "The effect of low level laser therapy on musculoskeletal pain: A meta-analysis". *Pain*. 1993; 52: 63-6.
  22. Olavi A, Pekka R, Pertti K, Pekka P. "Effects of the infrared laser therapy at treated and non-treated trigger points". *Acupunct. Electrother Res*. 1989; 14: 9-14.
  23. Thorsen H, Gam A, Swensson B, Jess M, Jensen M, Picullel I, et al. "Low level laser therapy for myofascial pain in the neck and shoulder girdle. A doubleblind, cross-over study". *Scand J. Rheumatol*. 1992; 21 (3): 139-141.
  24. Saygun I, Nizam N, Ural A, Sendar MA, Avcu F, Tozum TF. "Low-level laser irradiation affects the release of basic fibroblast growth factor (bFGF), insulin-like growth factor-I (IGF-I), and receptor of IGF-I (IGFBP3) from osteoblasts". *Photomed Laser Surg*. 2012; 30 (3): 149-154.
  25. Esmaeelinejad M, Bayat M. "Effect of low-level laser therapy on the release of interleukin-6 and basic fibroblast growth factor from cultured human skin fibroblasts in normal and high glucose mediums". *J Cosmet Laser Ther*. 2013; 15 (6): 310-317.
  26. Conroy DE, Hayes KW. "The effect of joint mobilization as a component of comprehensive treatment for primary shoulder impingement syndrome". *J Orthop Sports Phys Ther*. 1998; 28(1): 3-14.
  27. Michener LA, Walsworth MK, Burnet EN. "Effectiveness of rehabilitation for Patients with subacromial impingement syndrome: a systematic review". *J Hand Ther*. 2004; 17 (2): 152-164.
  28. Aktas I, Akgun K. and Cakmak B. "Therapeutic effect of pulsed electromagnetic field in conservative treatment of subacromial impingement syndrome". *Clin Rheumatol*. 2007; 26 (8): 1234-1239.
  29. Senbursa G, Baltaci G, Atay A. "Comparison of conservative treatment with and without manual physical therapy for patients with shoulder impingement syndrome: a prospective, randomized clinical trial". *Knee Surg Sports Traumatol Arthrosc*. 2007; 15(7): 915-921.
  30. Tate AR, McClure PW, Young IA, Salvatori R, Michener LA. "Comprehensive impairment-based exercise and manual therapy intervention for patients with subacromial impingement syndrome: a case series". *J Orthop Sports Phys Ther*. 2010; 40 (8): 474-493.